HIGHLIGHTS OF PRESCRIBING INFORMATION
These highlights do not include all the information needed to use PIMJETMI safely and effectively. See full prescribing information for PIMJETMI.

PIMJETMI® (pimicotinib) capsules, for oral use
Initial U.S. Approval: 2026

----------INDICATIONS AND USAGE----------

PIMJETMI is a kinase inhibitor indicated for the treatment of adult patients with symptomatic tenosynovial giant cell tumor (TGCT) for which surgical resection may cause worsening functional limitation or severe morbidity. (1)

----------DOSAGE AND ADMINISTRATION----------

- Recommended dosage: 50 mg orally once daily with or without food. (2.1)

----------DOSAGE FORMS AND STRENGTHS----------

Capsules: 25 mg of pimicotinib. (3)

----------CONTRAINDICATIONS----------

None. (4)

----------WARNINGS AND PRECAUTIONS----------

- Hepatotoxicity: Elevated AST and ALT can occur. Evaluate liver tests prior to initiation of treatment and during treatment. (5.1)
- Embryo-fetal toxicity: Can cause fetal harm. Advise of the potential risk to a fetus and to use effective contraception. (5.2, 8.1, 8.3)
- Increased serum creatinine without affecting renal function: Increases in serum creatinine can occur. Use alternative measures that are not based on serum creatinine to assess renal function. (5.3)

----------ADVERSE REACTIONS----------

The most common (>20%) adverse reactions were pruritus, rash, face edema, periorbital edema, fatigue, nausea, headache, and peripheral edema.

The most common (>30%) lab abnormalities were creatine phosphokinase increased, lactate dehydrogenase increased, aspartate aminotransferase increased, lipase increased, cholesterol increased, amylase increased, and alanine aminotransferase increased.

To report SUSPECTED ADVERSE REACTIONS, contact EMD Serono at 1-800-283-8088 ext. 5563 or FDA at 1-800-FDA-1088 or *www.fda.gov/medwatch*.

----------DRUG INTERACTIONS----------

- Strong or moderate CYP3A inhibitors: Avoid concomitant use with PIMJETMI. If concomitant use cannot be avoided, reduce the PIMJETMI dosage. (2.3, 7.1)
- Strong CYP3A inducers: Avoid concomitant use with PIMJETMI. (2.3, 7.1)
- Moderate CYP3A inducers: Avoid concomitant use with PIMJETMI. If concomitant use cannot be avoided, increase the PIMJETMI dosage. (2.3, 7.1)

----------USE IN SPECIFIC POPULATIONS----------

Lactation: Advise not to breastfeed. (8.2)

See 17 for PATIENT COUNSELING INFORMATION and FDA-approved Medication Guide.

Revised: 11/2026

FULL PRESCRIBING INFORMATION: CONTENTS*

1 INDICATIONS AND USAGE
2 DOSAGE AND ADMINISTRATION
2.1 Recommended Dosage and Administration
2.2 Dosage Modifications for Adverse Reactions
2.3 Dosage Modification for Drug Interactions
3 DOSAGE FORMS AND STRENGTHS
4 CONTRAINDICATIONS
5 WARNINGS AND PRECAUTIONS
5.1 Hepatotoxicity
5.2 Embryo-Fetal Toxicity
5.3 Increased Creatinine without Affecting Renal Function
6 ADVERSE REACTIONS
6.1 Clinical Trials Experience
7 DRUG INTERACTIONS
7.1 Effects of Other Drugs on PIMJETMI
8 USE IN SPECIFIC POPULATIONS
8.1 Pregnancy
8.2 Lactation
8.3 Females and Males of Reproductive Potential
8.4 Pediatric Use
8.5 Geriatric Use
11 DESCRIPTION
12 CLINICAL PHARMACOLOGY
12.1 Mechanism of Action
12.2 Pharmacodynamics
12.3 Pharmacokinetics
13 NONCLINICAL TOXICOLOGY
13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility
13.2 Animal Toxicology and/or Pharmacology
14 CLINICAL STUDIES
16 HOW SUPPLIED/STORAGE AND HANDLING
17 PATIENT COUNSELING INFORMATION

*Sections or subsections omitted from the full prescribing information are not listed

FULL PRESCRIBING INFORMATION

1 INDICATIONS AND USAGE

PIMJETMI is indicated for the treatment of adult patients with symptomatic tenosynovial giant cell tumor (TGCT) for which surgical resection may cause worsening functional limitations or severe morbidity.

2 DOSAGE AND ADMINISTRATION

2.1 Recommended Dosage and Administration

The recommended dosage of PIMJETMI is 50 mg orally once daily with or without food.

Swallow PIMJETMI capsules whole. Do not open, break, or chew the capsules.

Missed Dose

If a dose of PIMJETMI is missed within 12 hours, take the missed dose. If a dose is missed by more than 12 hours, resume at the next regularly scheduled time.

Vomiting

If vomiting occurs after taking a dose, do not take an additional dose and take the next dose at the regularly scheduled time.

2.2 Dosage Modifications for Adverse Reactions

Temporarily withhold PIMJETMI or reduce the dose to 25 mg once daily to manage adverse reactions *[see Adverse Reactions (6.1)]*.

Permanently discontinue PIMJETMI in patients who are unable to tolerate 25 mg orally daily.

The recommended dosage modifications of PIMJETMI for hepatotoxicity are provided in Table 1 *[see Warnings and Precautions (5.1)]*.

Table 1: Recommended Dosage Modifications for Hepatotoxicity

Hepatotoxicity Severity	**PIMJETMI Dosage Modifications**
AST or ALT increases >5 to 8 x ULN	• Withhold PIMJETMI and monitor weekly until recovery to < 3 times ULN or patient's baseline. • Resume PIMJETMI at a reduced dose of 25 mg once daily.
AST or ALT >3 to 5 × ULN and total bilirubin >2 × ULN	• Withhold PIMJETMI. • Evaluate patient for drug-induced liver injury or alternative causes. • If drug-induced liver injury confirmed, permanently discontinue PIMJETMI. • If drug-induced liver injury is ruled out, resume treatment at reduced dose of 25 mg once daily when ALT and AST decreased to <3 times ULN or patient's baseline and total bilirubin decreased to baseline. Monitor the patient.
AST or ALT >5 × ULN and total bilirubin >1.5 × ULN	• Permanently discontinue PIMJETMI.

AST or ALT >8 × ULN	• Permanently discontinue PIMJETMI.
Any AST or ALT and total bilirubin >3 × ULN	• Permanently discontinue PIMJETMI.

ALT - alanine aminotransferase; AST - aspartate aminotransferase; ULN - upper limit of normal

2.3 Dosage Modifications for Drug Interactions

Strong and Moderate CYP3A Inhibitors

Avoid concomitant use of strong and moderate CYP3A inhibitors.

If concomitant use cannot be avoided, reduce PIMJETMI dose from 50 mg once daily to 25 mg once daily *[see Drug Interactions (7.1)]*.

After the strong or moderate inhibitor has been discontinued for 3-5 half-lives, resume the PIMJETMI dosage taken prior to initiating the inhibitor.

Strong and Moderate CYP3A Inducers

Avoid concomitant use of strong and moderate CYP3A inducers.

If concomitant use of a moderate CYP3A inducer cannot be avoided, increase the PIMJETMI dosage from 50 mg once daily to 75 mg once daily *[see Drug Interactions (7.1)]*.

After discontinuing use of a moderate CYP3A inducer, resume PIMJETMI dose (7 to 14 days after discontinuing the CYP3A inducer) that was taken prior to initiating the moderate CYP3A inducer.

3 DOSAGE FORMS AND STRENGTHS

Capsules: 25 mg of pimicotinib, white opaque, size 3, imprinted with “PIMI 25” in black ink.

4 CONTRAINDICATIONS

None.

5 WARNINGS AND PRECAUTIONS

5.1 Hepatotoxicity

Cases of serious and fatal liver injury have occurred with the use of another kinase inhibitor that targets CSF1R *[see Clinical Pharmacology (12.1)]*. Serious and fatal liver injury have not been observed with PIMJETMI.

Across clinical trials in 255 patients treated with PIMJETMI, 1 (0.4%) patient had Grade 3 increased ALT and 2 (0.8%) patients had Grade 3 increased AST.

Dosage interruptions occurred in 3.5% of patients and dose reductions in 0.8% of patients due to AST/ALT increase.

Avoid PIMJETMI in patients with pre-existing active liver or biliary disease. Monitor liver function tests (including AST, ALT, ALP, gamma-glutamyl transferase (GGT) and total bilirubin), prior to initiation of PIMJETMI, monthly for the first 3 months then every 3 months for the first year of therapy and as clinically indicated thereafter. Withhold and reduce the dose, or permanently discontinue PIMJETMI based on the severity of liver transaminase elevations *[see Dosage and Administration (2.2)]*.

5.2 Embryo-Fetal Toxicity

Based on animal studies and its mechanism of action, PIMJETMI can cause fetal harm when administered to a pregnant woman. Oral administration of pimicotinib to pregnant rats and rabbits during organogenesis resulted in embryo-fetal toxicity including structural abnormalities and alterations to growth at exposures as low as 3 times the recommended dose based on area under the curve (AUC). Oral administration of pimicotinib to rats during organogenesis, parturition, and lactation resulted in neonatal mortality and alterations to growth at exposures below the recommended dose based on AUC. Advise pregnant women of the potential risk to a fetus. Advise females of reproductive potential and males with female partners of reproductive potential to use effective contraception during treatment with PIMJETMI and for 2 weeks after the last dose *[see Use in Specific Populations (8.1, 8.3)]*.

5.3 Increased Serum Creatinine Without Affecting Renal Function

In MANEUVER study Part 1, a maximum mean increase in serum creatinine of 11.4 µmol/L was observed 5 weeks after initiation of treatment with PIMJETMI. The serum creatinine increases persisted throughout treatment and were reversible upon treatment completion. These increases in serum creatinine may not be associated with changes in renal function. The increases in serum creatinine may be due to inhibition of renal tubular secretion transporters *[see Clinical Pharmacology (12.3)]*. During PIMJETMI treatment, use alternative measures that are not based on serum creatinine to assess renal function.

6 ADVERSE REACTIONS

The following clinically significant adverse reactions are described in greater detail elsewhere in the labeling:

- Hepatotoxicity *[see Warnings and Precautions (5.1)]*

6.1 Clinical Trials Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in practice.

The pooled safety population described in the Warnings and Precautions reflects exposure to PIMJETMI in 255 patients including 63 patients enrolled in the double-blind portion and in 31 patients in the open-label portion who crossed over to PIMJETMI in MANEUVER *[see Clinical Studies (14)]* and in 161 patients with TGCT or solid tumors in other clinical trials. The safety of PIMJETMI was evaluated in 63 adult patients with TGCT in MANEUVER *[see Clinical Studies (14)]*. MANEUVER excluded patients with bilirubin, AST, or ALT >1.5 x ULN. All patients received PIMJETMI 50 mg daily until disease progression or unacceptable toxicity. At completion of the double-blind, randomized part of the trial, 78% of patients treated with PIMJETMI were exposed for 6 months or longer and 37% were exposed for greater than one year.

Serious adverse reactions were reported in 3 (4.8%) patients who received PIMJETMI. Serious adverse reactions in ≥1% included erythema nodosum, infectious enterocolitis, and blood pressure increase.

Permanent discontinuation due to an adverse reaction occurred in 1 (1.6%) patient (fatigue) who received PIMJETMI.

Dose reductions due to an adverse reaction occurred in 8% of patients who received PIMJETMI. Adverse reactions leading to dose reduction in >1% of patients were rash, creatine phosphokinase increased, fatigue, and headache.

Treatment interruptions due to an adverse reaction occurred in 56% of patients who received PIMJETMI. Adverse reactions leading to dose interruption in >2% of patients were creatine phosphokinase increased,

rash, face edema, periorbital edema, pruritus, peripheral edema, neutrophil count decreased, AST increase, lipase increased, fatigue, and hypertension.

The most common (>20%) adverse reactions were pruritus, rash, face edema, periorbital edema, fatigue, nausea, headache, and peripheral edema.

The most common (>30%) lab abnormalities were creatine phosphokinase increased, lactate dehydrogenase increased, aspartate aminotransferase increased, lipase increased, cholesterol increased, amylase increased, and alanine aminotransferase increased.

Table 2 and Table 3 summarize the adverse reactions and laboratory abnormalities in MANEUVER during the randomized phase through Week 25.

Table 2: Adverse Reactions, Occurring in ≥10% of Patients Receiving PIMJETMI with a Difference Between Arms of >5% Compared to Placebo Through Week 25 in MANEUVER

System organ class Adverse reaction[#]	PIMJETMI N=63		Placebo N=31	
	All Grades %	Grade 3 or 4 %	All Grades %	Grade 3 or 4 %
Skin and subcutaneous tissue disorders				
Pruritus	52	3.2	3.2	0
Rash[a]	48	6	10	0
General disorders and administration site conditions				
Face edema	48	0	0	0
Fatigue[a]	37	0	23	0
Peripheral edema[a]	21	0	6	0
Eye disorders				
Periorbital edema[a]	41	0	10	0
Gastrointestinal disorders				
Nausea	27	0	6	0
Nervous system disorders				
Headache[a]	22	0	6	0
Dizziness	16	0	3.2	0
Vascular disorders				
Hypertension[a]	19	10	6	0
Hemorrhage[a]	13	0	0	0
Psychiatric disorders				
Insomnia	14	0	6	0

The severity of adverse reactions was assessed using CTCAE v5.0.

[a] Includes multiple related terms

Table 3: Laboratory Abnormalities Worsening from Baseline in ≥10% of Patients Receiving PIMJETMI with a Difference Between Arms of >5% Compared to Placebo Through Week 25 in MANEUVER

Laboratory Abnormality[a, #]	PIMJETMI N=63		Placebo N=31	
	All Grades %	Grade 3 or 4 %	All Grades %	Grade 3 or 4 %
Chemistry				
CPK increased	97	16*	19	0
LDH increased	95	0	6	0
AST increased	83	0	13	0
Lipase increased	58	4.2	26	0
Cholesterol increased	53	0	30	0
Amylase increased	52	0	3.3	0
ALT increased	37	0	26	0
Creatinine increased	25	0	0	0
Hematology				
White blood cell decreased	27	0	3.2	0
Neutrophil count decreased	21	1.6	0	0
Lymphocyte count decreased	14	0	3.2	0

ALT - alanine aminotransferase; AST - aspartate aminotransferase; CPK - creatine phosphokinase; LDH - blood lactate dehydrogenase

The severity of adverse reactions was assessed using CTCAE v5.0.

* Includes Grade 4 events

[a] The denominator used to calculate the rate varied from 48 to 63 for PIMJETMI and from 23 to 31 for Placebo based on the number of patients with a baseline value and at least one post-treatment value. Represents the incidence of laboratory findings, not of reported adverse events

Other Clinical Trial Experience

Other clinically significant adverse reactions occurring across clinical trials in patients with TGCT treated with PIMJETMI 50 mg once daily (N = 158) include dermal mucinosis reported as skin indentations (23%).

7 DRUG INTERACTIONS

7.1 Effects of Other Drugs on PIMJETMI

Table 4 describes drug interactions where concomitant use of another drug affects PIMJETMI.

Table 4: Effects of Other Drugs on PIMJETMI

Strong and Moderate CYP3A Inhibitors	
Prevention or management	• Avoid concomitant use of strong and moderate CYP3A inhibitors with PIMJETMI. • If concomitant use cannot be avoided, reduce the PIMJETMI dosage *[see Dosage and Administration (2.3)]*.
Mechanism and Clinical Effects	Pimicotinib is a CYP3A substrate. Concomitant use with a strong or moderate CYP3A inhibitor increases pimicotinib exposure *[see Clinical Pharmacology (12.3)]*, which may increase the risk of PIMJETMI-related adverse reactions.
Strong and Moderate CYP3A Inducers	
Prevention or management	• Avoid concomitant use of strong and moderate CYP3A inducers with PIMJETMI. • If concomitant use cannot be avoided with a moderate CYP3A inducer, increase the PIMJETMI dosage *[see Dosage and Administration (2.3)]*.
Mechanism and Clinical Effects	Pimicotinib is a CYP3A substrate. Concomitant use with a strong or moderate CYP3A inducer decreases pimicotinib exposure *[see Clinical Pharmacology (12.3)]*, which may reduce the effectiveness of PIMJETMI.

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Risk Summary

Based on animal studies and its mechanism of action, PIMJETMI can cause fetal harm when administered to a pregnant woman *[see Clinical Pharmacology (12.1)]*. There are no available data on the use of PIMJETMI in pregnant women.

Oral administration of pimicotinib to pregnant rats and rabbits during organogenesis resulted in embryo-fetal toxicity including structural abnormalities and alterations to growth at exposures as low as 3 times the recommended dose based on AUC. Oral administration of pimicotinib to rats during organogenesis, parturition, and lactation resulted in neonatal mortality and alterations to growth at exposures below the recommended dose based on AUC (*see Data*). Advise pregnant women of the potential risk to a fetus.

In the U.S. general population, the estimated background risk of major birth defects and miscarriage in clinically recognized pregnancies is 2% to 4% and 15% to 20%, respectively.

Data

Animal Data

In embryo-fetal development studies, pimicotinib was orally administered to rats and rabbits during the period of organogenesis. In rats, structural abnormalities including external malformation and subcutaneous edema were observed at a dose of 10 mg/kg/day (18 times the exposure at the recommended dose based on AUC). In rabbits, structural abnormalities, including external and visceral malformations, were observed at doses as low as 5 mg/kg/day (3 times the exposure at the recommended

dose based on AUC). Fetal growth restriction, including decreased fetal weight and crown-rump length, and increased visceral variations were observed at 20 mg/kg/day (15 times the exposure at the recommended dose based on AUC).

In a pre-and postnatal developmental study in rats, pimicotinib was orally administered to rats at doses of 1, 2, and 4 mg/kg/day during organogenesis, parturition, and lactation. Maternal toxicity including mortality was observed at 2 mg/kg/day. Fetal growth restriction and neonatal mortality were observed at ≥1 mg/kg/day (0.8 times the exposure at the recommended dose based on AUC).

8.2 Lactation

Risk Summary

There are no data on the presence of pimicotinib or its metabolites in human milk or its effects on a breastfed child or on milk production. When administered to lactating rats, pimicotinib was detected in the plasma of nursing pups (*see Data*). When a drug is present in animal milk, it will likely be present in human milk. Because of the potential for serious adverse reactions in a breastfed child, advise women not to breastfeed during treatment with PIMJETMI and for 2 weeks after the last dose.

Data

Animal Data

In a pre- and postnatal developmental study in rats, pimicotinib was orally administered to lactating rats at doses of 1, 2, and 4 mg/kg/day. Pimicotinib was detected in the plasma of nursing pups on postnatal days 4 and 21, indicating the presence of pimicotinib in rat milk or prolonged persistence in pups.

8.3 Females and Males of Reproductive Potential

PIMJETMI can cause fetal harm when administered to a pregnant woman *[see Use in Specific Populations (8.1)]*.

Pregnancy Testing

Verify pregnancy status of females of reproductive potential prior to initiating treatment with PIMJETMI.

Contraception

Females

Advise females of reproductive potential to use effective contraception during treatment with PIMJETMI and for 2 weeks after the last dose.

Males

Advise males with female partners of reproductive potential to use effective contraception during treatment with PIMJETMI and for 2 weeks after the last dose.

Infertility

Males

Based on findings in animals, PIMJETMI may impair male fertility. The effect on sperm in rats was reversible *[see Nonclinical Toxicology (13.1)]*.

8.4 Pediatric Use

Safety and effectiveness of PIMJETMI have not been established in pediatric patients.

Animal Toxicity Data

In a rat repeat-dose toxicology study, oral pimicotinib resulted in decreased systemic calcium and phosphate levels and increased thickness of the physis and trabecular bone in the distal femur and proximal tibia at ≥3 mg/kg (7 times the exposure at the recommended dose based on AUC).


8.5 Geriatric Use

Of the 63 patients with TGCT treated with PIMJETMI in the MANEUVER trial, 6 (10%) were 65 years of age and older, while none were 75 years of age and older; this trial did not include sufficient numbers of patients aged 65 and older to determine whether they respond differently from younger patients.

11 DESCRIPTION

Pimicotinib is a kinase inhibitor. The chemical name of pimicotinib hydrochloride is 3,3-dimethyl-*N*-(6-methyl-5-{[2-(1-methyl-1*H*-pyrazol-4-yl)pyridin-4-yl]oxy}pyridin-2-yl)-2-oxopyrrolidine-1-carboxamide hydrochloride hydrate. The molecular formula for pimicotinib hydrochloride is $C_{22}H_{24}N_6O_3$•HCl•H_2O. The molecular weight is 474.94 g/mol for the pimicotinib hydrochloride and 420.47 g/mol for the free base.

The chemical structure is shown below.



Pimicotinib hydrochloride is a white to off-white powder. The solubility of pimicotinib hydrochloride in aqueous solutions and bio-relevant media decreases with increasing pH. The pKa was determined to be 5.0 (base). Pimicotinib hydrochloride is freely soluble in ethanol and practically insoluble or insoluble in water.

PIMJETMI (pimicotinib) capsules for oral administration contain 25 mg pimicotinib equivalent to 28.24 mg pimicotinib hydrochloride monohydrate. Each capsule contains the following inactive ingredients: colloidal silicon dioxide, croscarmellose sodium, lactose monohydrate, magnesium stearate, and microcrystalline cellulose. The capsule shell, imprinted with black ink, contains the following inactive ingredients: black iron oxide, butyl alcohol, dehydrated alcohol, gelatin, isopropyl alcohol, potassium hydroxide, propylene glycol, purified water, shellac, strong ammonia solution, and titanium dioxide.

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

Pimicotinib is a kinase inhibitor that inhibits colony stimulating factor 1 receptor (CSF-1R). In vitro, pimicotinib inhibited CSF-1R kinase activity and CSF-1R-dependent cell proliferation.

12.2 Pharmacodynamics

Exposure-Response Relationship

Higher pimicotinib exposure is associated with an increased risk of elevated amylase and elevated AST.

Pimicotinib exposure-response relationship and time course of pharmacodynamic response for the effectiveness have not been fully characterized.

Cardiac Electrophysiology

At the maximum recommended PIMJETMI dosage (50 mg once daily), a mean increase in the QTc interval >20 ms is unlikely.

12.3 Pharmacokinetics

Pimicotinib pharmacokinetics were observed at steady state in patients with advanced solid tumors at the approved recommended dosage and are presented as mean (coefficient of variation, %), unless otherwise specified. Pimicotinib maximum concentration (C_{max}) is 566 (66%) ng/mL and total systemic exposure (AUC) is 5,692 (62%) hrs*ng/mL. C_{max} and AUC increased in a dose proportional manner over the dose range from 25 to 75 mg (0.5 to 1.5 times the recommended dose). Steady state is reached within 15 days and the mean accumulation ratio based on AUC is 3.4.

Absorption

Pimicotinib median (min, max) time to maximum plasma concentrations (t_{max}) is 1 (0.4, 3.5) hour.

Effect of Food

Pimicotinib C_{max} decreased to 40% with no change in AUC following administration of a high-fat, high-calorie meal (total 800-1,000 calories, of which 500-600 from fat).

Distribution

Pimicotinib apparent (oral) volume of distribution is 730 L (38%).

Pimicotinib plasma protein binding is 95% and is not concentration dependent, in vitro.

Pimicotinib blood-to-plasma ratio is about 0.8 to 1.3 in humans.

Elimination

Pimicotinib elimination half-life is 44 hours (34%) with an apparent (oral) clearance of 12 L/h (53%).

Metabolism

The metabolic pathways of pimicotinib include primarily hydrolysis and oxidation. Oxidative metabolism is mediated mainly via CYP3A and accounts for approximately 20% of total drug elimination.

Excretion

Following a single dose of radiolabeled pimicotinib 50 mg to healthy subjects, 58% of the dose was recovered in urine and 38% in feces with no notable unchanged pimicotinib in urine or feces.

Specific Populations

No clinically significant differences in the pharmacokinetics of pimicotinib were observed based on age (18 to 82 years), body weight (41 to 120 kg), sex, race (73% Asian, 5% Black or African American or 21% White), or CLcr 30 to 89 mL/min (estimated by Cockcroft-Gault). The effect of CLcr < 30 mL/min and dialysis on the pharmacokinetics of pimicotinib have not been studied.

Patients with Hepatic Impairment

No clinically significant changes in the pharmacokinetics of pimicotinib were observed in patients with mild hepatic impairment (Child Pugh class A). Pimicotinib AUC increased 1.5-fold in subjects with moderate hepatic impairment (Child Pugh class B) with no change in C_{max}. The effect of severe hepatic impairment (Child Pugh class C) has not been studied.

Drug Interaction Studies

Clinical Studies and Model-Informed Approaches

Strong and Moderate CYP3A Inhibitors: Pimicotinib AUC increased 1.6-fold with no change in C_{max} following concomitant use of itraconazole (strong CYP3A4 inhibitor) 200 mg once daily for 4 days.

Pimicotinib AUC is predicted to increase 1.4-fold to 1.5-fold with no change in C_{max} following concomitant use of fluconazole or erythromycin (moderate CYP3A4 inhibitors).

Strong and Moderate CYP3A Inducers: Pimicotinib AUC decreased to 31% and C_{max} to 72% following concomitant use of rifampicin (strong CYP3A inducer) 600 mg QD for 7 days.

Pimicotinib AUC is predicted to decrease to 48% and C_{max} to 78% following concomitant use of efavirenz (moderate CYP3A inducer).

Acid-Reducing Agents: Pimicotinib displayed pH-dependent aqueous solubility *[see Description (11)]*. Pimicotinib C_{max} decreased to 58% with no change in AUC following concomitant use of omeprazole 40 mg QD for 5 days.

CYP3A Substrates: Midazolam (CYP3A substrate) AUC and C_{max} are predicted to increase 1.4-fold following concomitant use of multiple doses of PIMJETMI.

P-gp Substrates: Fexofenadine (P-gp substrate) AUC increased 1.6-fold and C_{max} 1.8-fold following concomitant use of multiple doses of PIMJETMI.

MATE1 Substrates: Metformin (MATE1 substrate) AUC increased 1.4-fold and C_{max} 1.3-fold following concomitant use of multiple doses of PIMJETMI.

BCRP Substrates: Rosuvastatin (BCRP substrate) AUC increased 1.4-fold and C_{max} 1.5-fold following concomitant use of multiple doses of PIMJETMI.

In Vitro Studies

CYP enzymes: Pimicotinib does not inhibit CYP1A2, CYP2B6, CYP2C9, CYP2C19, CYP2C8 or CYP2D6. Pimicotinib does not induce CYP1A2 or CYP2B6.

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

Carcinogenesis

Pimicotinib was not carcinogenic in a 6-month rasH2-Tg mouse study at oral doses up to 40 mg/kg/day (approximately 50 times the exposure at the recommended dose based on AUC).

Mutagenesis

Pimicotinib and its major human circulating metabolites, M281 and M113, were not mutagenic in the bacterial reverse mutation test (Ames) or clastogenic in an in vitro chromosomal aberration assay. Pimicotinib was not clastogenic or aneugenic in an in vivo bone marrow micronucleus assay in mice at doses up to 200 mg/kg/day. M113 was not clastogenic or aneugenic in an in vivo bone marrow micronucleus assay in rats at doses up to 1000 mg/kg/day in males or 750 mg/kg/day in females.

Impairment of Fertility

In a fertility and early embryonic development study, no effects on fertility or reproductive capacities were observed in male or female rats at oral pimicotinib doses up to 20 mg/kg/day (37 times the exposure at the recommended dose based on AUC). In a 26-week rat repeat-dose toxicology study, sperm were decreased at ≥3 mg/kg/day (7 times the exposure at the recommended dose based on AUC), but the finding was reversible.

13.2 Animal Toxicology and/or Pharmacology

In a rat repeat-dose toxicology study, pimicotinib resulted in glomerulonephritis and degeneration of blood vessels (vascular inflammation and necrosis of arteries) at ≥1 mg/kg/day (approximately 2 times the exposure at the recommended dose based on AUC).

14 CLINICAL STUDIES

The efficacy of PIMJETMI was evaluated in MANEUVER (NCT05804045), a double-blind, randomized (2:1), placebo-controlled, international, multicenter trial in patients with TGCT for whom surgical resection may cause dysfunction or serious complications. Eligible patients were symptomatic and had measurable disease per Response Evaluation Criteria in Solid Tumors (RECIST v1.1) with at least one lesion having a minimum size of 2 cm.

Patients were randomized to PIMJETMI 50 mg or placebo once daily. Randomization was stratified by geographic region (China versus non-China). At Week 25, patients who completed the double-blind, randomized part of the trial were eligible to participate in an open-label extension study in which all patients received PIMJETMI.

The major efficacy outcome measure was overall response rate (ORR), as assessed by blinded independent central review (BICR) per RECIST v1.1 at Week 25. Additional efficacy outcomes measured at Week 25 were ORR as assessed by BICR using tumor volume score (TVS), mean change from baseline in the range of motion (ROM) of the affected joint measured by goniometry assessments, Brief Pain Inventory (BPI) Worst Pain numeric rating scale (NRS) score, and Patient-reported Outcomes Measurement Information System Physical Functioning (PROMIS-PF) Short Form.

A total of 94 patients were randomized to PIMJETMI (N=63) or placebo (N=31) during the double-blind period of the study. The median age was 40 years (range: 18-69 years); 68% of patients were female; 51% were Asian, 39% were White, 5% were Black or African American, and 4.3% were Other; 87% were not Hispanic or Latino, 3.2% were Hispanic, and 10% were not reported or unknown; 85% of patients had diffuse TGCT; 60% had prior surgery, and 6% had previously been treated with systemic therapy. Disease locations were knee (50%), ankle (15%), hip (14%), foot (9%), wrist (7%), and other (5%).

A statistically significant improvement in ORR was demonstrated in patients randomized to PIMJETMI compared with placebo. Efficacy results in MANEUVER are summarized in Table 5.

Table 5: Efficacy Results Assessed at Week 25 for MANEUVER

Efficacy Parameter	**PIMJETMI N=63**	**Placebo N=31**
Overall response rate[a, b] (ORR) per RECIST v1.1		
ORR, n (%) (95% CI)	35 (56) (42, 68)	1 (3.2) (0.1, 17)
Complete response	1 (1.6)	0
Partial response	34 (54)	1 (3.2)
p-value	<0.0001	
Duration of response (DoR)[b]		
Median (months)[c] Range (months)	NR (2.5, 26.2+)	NA
Patients with DoR ≥ 6 months[d]	32/35 (91%)	-

Patients with DoR ≥ 12 months[d]	27/35 (77%)	-
Range of Motion (ROM)[e] (%)		
Patients with data at baseline and Week 25, n	55	28
at baseline, mean (SD)	70.6 (20.7)	66.1 (21.8)
at Week 25, mean (SD)	84.9 (21.4)	67.6 (23.6)
Change from baseline, LS mean (95% CI)[f]	15.6 (10.4, 20.9)	-0.1 (-7.0, 6.9)
Difference from placebo (95% CI)	15.7 (7.3, 24.1)	
p-value	0.0003	
BPI Worst Pain NRS[e, g]		
Patients with data at baseline and Week 25, n	56	29
at baseline, mean (SD)	4.8 (2.1)	4.8 (2.6)
at Week 25, mean (SD)	2.5 (1.9)	4.4 (2.5)
Change from baseline, LS mean (95% CI)[f]	-2.3 (-2.7, -1.9)	-0.2 (-0.8, 0.3)
Difference from placebo (95% CI)	-2.1 (-2.8, -1.4)	
p-value	<0.0001	
PROMIS Physical Function T-Score[e, h]		
Patients with data at baseline and Week 25, n	56	29
at baseline, mean (SD)	41.2 (7.1)	39.8 (6.0)
at Week 25, mean (SD)	46.6 (7.8)	42.6 (7.8)
Change from baseline, LS mean (95% CI)[f]	5.6 (4.2, 7.1)	2.2 (0.2, 4.2)
Difference from placebo (95% CI)	3.4 (0.9, 5.9)	
p-value	0.0074	

+: Indicated that the patient's response was ongoing at last assessment as of the data cutoff date
NR - Not reached; NA - Not applicable; BPI - Brief Pain Inventory; CI - confidence interval; LS - least squares; MMRM - mixed model for repeated measures; N - number of patients in treatment group in analysis set and denominators for all percentages; n - number of patients in analysis with data available; NRS - numerical rating scale; PROMIS-PF - Patient-reported Outcomes Measurement Information System- Physical Function; ROM - range of motion; SD - Standard deviation.

[a] Assessed by blinded independent central review.

[b] ORR analysis was based on data cut-off date (DCO) March 12, 2025.

DOR results are based on an approximately additional 12-months of follow-up from the time of ORR analysis. DCO: March 1, 2026.

[c] Estimate using Kaplan-Meier method.

[d] Based on observed duration of response.

[e] ROM, BPI Worst Pain NRS and PROMIS PF T-Score endpoints analysis were based on DCO September 23, 2024.

[f] Estimated using a MMRM approach.


[g] Measured by BPI Worst Pain Numeric Rating Scale (NRS), based on a scale of 0 to 10. Low numbers on scale indicate less pain.

[h] Measured by PROMIS-PF T-score (15-Item score; ranges from 0-100). Higher scores of PROMIS-PF indicate better physical functioning.

ORR per RECIST v1.1 by BICR for patients receiving PIMJETMI in China (N=31) was 65% (95% CI: 45, 81) and 47% (95% CI: 29, 65) outside of China (N=32). ORR as assessed by the Investigator was 35% (95% CI: 23%, 48%) in the PIMJETMI arm and 0% in the placebo arm. ORR by TVS was 65% (95% CI: 52%, 77%; $p < 0.0001$) in patients receiving PIMJETMI and 3.2% (95% CI: 0.1%,17%) in patients receiving placebo (DCO: March 12, 2025).

Figure 1 shows the change from baseline in range of motion for each patient at Week 25.

Figure 1: Change from Baseline in Range of Motion at Week 25 for MANEUVER



*Percent normal reference range for the affected joint.
+ Response by RECIST

16 HOW SUPPLIED/STORAGE AND HANDLING

PIMJETMI (pimicotinib) capsules are white opaque, size 3, imprinted with “PIMI 25” in black ink and contain 25 mg pimicotinib. Available in a carton (NDC 44087-6100-6) containing 2 bottles fitted with a child-resistant closure each containing 30 capsules and a desiccant (NDC 44087-6100-3).

Store PIMJETMI at 20ºC to 25ºC (68ºF to 77ºF); excursions permitted from 15ºC to 30ºC (59ºF to 86ºF) *[see USP Controlled Room Temperature]*.

Store in the original bottle with desiccant to protect from moisture.

17 PATIENT COUNSELING INFORMATION

Advise the patient to read the FDA-approved patient labeling (Medication Guide).

Hepatotoxicity

Advise patients that PIMJETMI can cause asymptomatic AST or ALT elevations and that their healthcare provider needs to monitor liver transaminase levels before and during initial few months of treatment *[see Warnings and Precautions (5.1)]*.

Embryo-Fetal Toxicity

Advise pregnant women and patients of reproductive potential of the potential risk to a fetus. Advise females of reproductive potential to inform their healthcare provider of a known or suspected pregnancy *[see Warnings and Precautions (5.2), Use in Specific Populations (8.1)]*.

Advise females of reproductive potential to use effective contraception during treatment with PIMJETMI and for 2 weeks after the last dose *[see Use in Specific Populations (8.3)]*.

Advise male patients with female partners of reproductive potential to use effective contraception during treatment with PIMJETMI and for 2 weeks after the last dose *[see Use in Specific Populations (8.3)]*.

Lactation

Advise females not to breastfeed during treatment with PIMJETMI and for 2 weeks after the last dose *[see Use in Specific Populations (8.2)]*.

Infertility

Advise male patients that PIMJETMI may impair fertility *[see Use in Specific Populations (8.3), Nonclinical Toxicology (13.1)]*.

Drug Interactions

Advise patients to inform their healthcare provider of all concomitant medications, including prescription medicines, over-the-counter drugs and herbal products *[see Drug Interactions (7)]*.

Manufactured for: EMD Serono, Inc., Boston, MA 02210, U.S.A.
PIMJETMI is a trademark of Merck KGaA, Darmstadt, Germany

MEDICATION GUIDE

PIMJETMI® (pim-JET-mee)

(pimicotinib)

capsules

What is the most important information I should know about PIMJETMI?

PIMJETMI can cause serious side effects, including:

Liver problems. Increased liver enzymes in your blood are common with PIMJETMI. Your healthcare provider will do blood tests to check for liver problems:

- before starting treatment with PIMJETMI,
- every month for the first 3 months,
- then every 3 months for the rest of the first year after starting treatment and as needed thereafter during treatment with PIMJETMI.

If you develop liver problems during treatment with PIMJETMI, your healthcare provider may temporarily stop treatment, decrease your dose, or permanently stop treatment depending on how severe your liver problems are.

Tell your healthcare provider right away if you develop any signs or symptoms of liver problems during treatment with PIMJETMI:

- yellowing of your skin or the white part of your eyes
- dark urine
- lack or loss of appetite
- pain or tenderness on the upper right side of your stomach-area (abdomen)
- feeling overly tired
- nausea
- vomiting
- fever
- rash
- itching

See "What are the possible side effects of PIMJETMI?" for information about side effects.

What is PIMJETMI?

PIMJETMI is a prescription medicine used to treat adults with symptomatic tenosynovial giant cell tumor (TGCT) when surgery may make the symptoms worse or cause severe problems.

It is not known if PIMJETMI is safe and effective in children.

Before taking PIMJETMI, tell your healthcare provider about all of your medical conditions, including if you:

- have or had liver problems.
- are pregnant or plan to become pregnant. PIMJETMI can harm your unborn baby.
  Females who are able to become pregnant:
  - Your healthcare provider will do a pregnancy test before starting treatment with PIMJETMI.
  - Use effective birth control (contraception) during treatment and for 2 weeks after your last dose of PIMJETMI.
  - Tell your healthcare provider right away if you become pregnant or you think you may be pregnant during treatment with PIMJETMI.

  Males with female partners who are able to become pregnant:
  - Use effective birth control (contraception) during treatment and for 2 weeks after your last dose of PIMJETMI.
- are breastfeeding or plan to breastfeed. It is not known if PIMJETMI passes into your breast milk. **Do not** breastfeed during treatment with PIMJETMI and for 2 weeks after your last dose of PIMJETMI.

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

Taking PIMJETMI with certain other medicines may affect how PIMJETMI works and can cause side effects.

How should I take PIMJETMI?

- Take PIMJETMI exactly as your healthcare provider tells you to take it.
- Take PIMJETMI 1 time a day.
- Take PIMJETMI with food or without food.
- Swallow PIMJETMI capsules whole. **Do not** open, break, or chew the capsules.
- If you miss a dose of PIMJETMI within 12 hours of your scheduled time, take the missed dose as soon as possible. If you miss your dose by more than 12 hours, skip the missed dose and take your next dose at your regularly scheduled time.
- If you vomit after taking a dose of PIMJETMI, do not take another dose and take the next dose at your regularly scheduled time.

What are the possible side effects of PIMJETMI?
PIMJETMI can cause serious side effects.
See "What is the most important information I should know about PIMJETMI?"

The most common side effects of PIMJETMI include:

- itching
- rash
- swelling of your face
- swelling around your eyes
- tiredness
- nausea
- headache
- swelling of your hands and feet
- change in certain blood tests

PIMJETMI may affect fertility in males, which may affect your ability to have a child. Talk to your healthcare provider if this is a concern for you.

These are not all of the possible side effects of PIMJETMI.
Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

How should I store PIMJETMI?

- Store PIMJETMI at room temperature between 68°F to 77° F (20°C to 25°C).
- PIMJETMI comes in a bottle with a child-resistant closure.
- PIMJETMI comes with a drying agent (desiccant) in the bottle. Store PIMJETMI in the original bottle with desiccant.
- Keep the bottle closed to protect from moisture.

Keep PIMJETMI and all medicines out of the reach of children.

General information about the safe and effective use of PIMJETMI

Medicines are sometimes prescribed for purposes other than those listed in a Patient Information leaflet. Do not use PIMJETMI for a condition for which it was not prescribed. Do not give PIMJETMI to other people, even if they have the same symptoms you have. It may harm them. You can ask your pharmacist or healthcare provider for information about PIMJETMI that is written for health professionals.

What are the ingredients in PIMJETMI?

Active ingredient: pimicotinib hydrochloride

Inactive ingredient: colloidal silicon dioxide, croscarmellose sodium, lactose monohydrate, magnesium stearate, and microcrystalline cellulose. The capsule shell, imprinted with black ink, contains the following inactive ingredients: black iron oxide, butyl alcohol, dehydrated alcohol, gelatin, isopropyl alcohol, potassium hydroxide, propylene glycol, purified water, shellac, strong ammonia solution, and titanium dioxide.

Manufactured for: EMD Serono, Inc, Boston, MA 02210, U.S.A.
PIMJETMI is a trademark of Merck KGaA, Darmstadt, Germany.
For more information, call 1-844-826-8371 or go to www.PIMJETMI.com.

This Patient Information has been approved by the U.S. Food and Drug Administration. Issued 11/2026